

THE HEALTHY MINDS STUDY

2014 DATA REPORT

ABOUT THE HEALTHY MINDS STUDY (HMS)



STUDY TEAM

Principal Investigator: Daniel Eisenberg, PhD

Co-Investigator: Sarah Ketchen Lipson, EdM

Research Study Coordinators: Katie Beck, Mira Dalal and Odessa Despot, PsyD

SURVEY SCIENCES GROUP, LLC

In 2013-2014 HMS was conducted in collaboration with the Survey Sciences Group, a survey research firm in Ann Arbor, Michigan. <http://www.surveysciences.com/>

GRAPHIC/LAYOUT DESIGNER

Sarah Fogel
University of Michigan School of Art and Design, Class of 2014

REPORT AUTOMATION

Andy Inscore
Software Engineer

TABLE OF CONTENTS

STUDY PURPOSE	1
STUDY DESIGN	2
ABOUT THIS REPORT	3
KEY FINDINGS	4
SAMPLE CHARACTERISTICS	5
PREVALENCE OF MENTAL HEALTH PROBLEMS	6
POSITIVE MENTAL HEALTH	7
HEALTH BEHAVIORS AND LIFESTYLE	8
ATTITUDES AND BELIEFS ABOUT MENTAL HEALTH SERVICES	9
USE OF SERVICES	10
ACADEMIC AND SOCIAL ENVIRONMENT	12
REFERENCES	14
APPENDIX: DESCRIPTIVE STATISTICS FOR ALL SURVEY ITEMS	15

STUDY PURPOSE

The Healthy Minds Study provides a detailed picture of mental health and related issues in college student populations. Schools typically use their data for some combination of the following purposes: to identify needs and priorities; benchmark against peer institutions; evaluate programs and policies; plan for services and programs; and advocate for resources.

STUDY DESIGN

The Healthy Minds Study is designed to protect the privacy and confidentiality of participants. HMS is approved by the Health Sciences and Behavioral Sciences Institutional Review Board at University of Michigan. To further protect respondent privacy, the study is covered by a Certificate of Confidentiality from the National Institutes of Health.

SAMPLING

Each participating school provides the HMS team with a randomly selected sample of up to 4,000 currently enrolled students over the age of 18. Schools with graduate students typically include both undergraduates and graduate students in the sample.

DATA COLLECTION

The HMS is a web-based survey. Students are invited and reminded to participate in the survey via emails, which are timed to avoid, if at all possible, the first two weeks of the term, the last week of the term, and any major holidays. The data collection protocol begins with an email invitation, and non-responders are contacted up to three times by email reminders spaced by 2-4 days each. Reminders are only sent to those who have not yet completed the survey. Each communication contains a URL and a unique study ID that students use to gain access to the survey. The HMS questionnaire asks up to 200 questions, with skip logic used to eliminate irrelevant questions.

NON-RESPONSE ANALYSIS

A potential concern in any survey study is that those who respond to the survey will not be fully representative of the population from which they are drawn. In the HMS, we can be confident that those who are invited to fill out the survey are representative of the full student population because these students are randomly selected from the full list of currently enrolled students. However it is still possible that those who actually complete the survey are different in important ways from those who do not complete the survey. The overall participation rate for the 2014 study was 29%. It is important to raise the question of whether the 29% who participated are different in important ways from the 71% who did not participate. We address this issue by constructing non-response weights using administrative data on full student populations. Nearly all of the 20 schools in the 2014 HMS were able to provide administrative data about all randomly selected students. The analysis of these administrative data, separated from any identifying information, was approved in the IRB application at the University of Michigan and at each participating school. We used the following variables, where available, to estimate which types of students were more or less likely to respond: gender, race/ethnicity, academic level, and grade point average. We used these variables to estimate the response propensity of each type of student (based on multivariate logistic regressions), and then assigned response propensity weights to each student who completed the survey. The less likely a type of student was to complete the survey, the larger the weight they received in the analysis, such that the weighted estimates are representative of the full student population in terms of the administrative variables available for each institution. Finally, note that these sample weights give equal aggregate weight to each school in the national estimates. An alternative would have been to assign weights in proportion to school size, but we decided that we did not want our overall national estimates to be dominated by schools in our sample with very large enrollments.

ABOUT THIS REPORT

This data report provides descriptive statistics (percentages, mean values, etc.) for a set of key measures using the aggregate sample of respondents across all colleges and universities that participated during the 2013-2014 academic year. In addition to the key measures highlighted in this report, an appendix is also included with descriptive statistics for each survey item (see below).

APPENDIX

The appendix includes values for most measures in the survey. For each measure, the table displays the following information: the national value (for all students, for male students, for female students) and the 95% confidence interval for the national value. All values in the appendix have been weighted to be representative of the full student populations to which they refer (see Non-response Analysis). Also note that for some measures, respondents were allowed to check more than one response category (e.g., they might have gone to more than one type of provider for mental health services), so the proportions sometimes add up to more than 1.00 across response categories. The 95% confidence intervals give a sense of how much uncertainty there is about each estimated value. This uncertainty exists because our estimates are based only on a random sample of students, rather than a complete census of the student population. However, some schools that had less than 4,000 students (the typical requested sample size), provided their entire population. For consistency sake, these schools were not treated any differently than those schools that provided a 4,000 student sample of their full population. Essentially, the confidence interval tells us that there is a 95% probability that the true population value is within this particular range.

EXPLORING THE DATA FURTHER

If you are interested in exploring the data beyond what is in this report, you can use a user-friendly website with drop-down menus, at data.healthymindsnetwork.org, or email us at healthyminds@umich.edu to request the full national data sets.


KEY FINDINGS

This section offers a quick look at results that may be of special interest to your institution.

Estimated values of selected measures for null	Proportion of students
Major depression (positive PHQ-9 screen)	12%
Depression overall, including major and moderate (positive PHQ-9 screen)	21%
Anxiety disorder (positive GAD-7screen)	22%
Eating disorder (positive SCOFF screen)	7%
Non-suicidal self-injury (past year)	20%
Suicidal ideation (past year)	10%
Lifetime diagnoses of mental disorders	34%
Psychiatric medication (past year)	18%
Mental health therapy/counseling (past year)	22%
Any mental health therapy/counseling and/or psychiatric medication among students with positive depression or anxiety screens (past year)	46%
Personal stigma: agrees with "I would think less of someone who has received mental health treatment."	7%
Perceived public stigma: agrees with "Most people would think less of someone who has received mental health treatment."	47%

SAMPLE CHARACTERISTICS (N=16342)

Age(years)




Gender




55% Female
45% Male
1% Transgender

Race/ethnicity




WHI White or Caucasian
BLA African American/Black
LAT Hispanic/Latino
AMIN American Indian/Alaskan Native
ARAB Arab/Middle Eastern or Arab American
ASIAN Asian/Asian-American
PAC Pacific Islander
OTH Other (specify)
N/A Not applicable - I would prefer not to identify my race/ethnicity

Living arrangement



27% Campus Residence Hall
1% Fraternity or sorority house
4% Other university housing
52% Off-campus, non-university housing
14% Parent or guardian's home
2% Other

Degree program




ASS Associate's degree
BA Bachelor's degree
MA Master's degree
JD JD
MD MD
PHD PhD or equivalent
OTH Other (specify)

PREVALENCE OF MENTAL HEALTH PROBLEMS


DEPRESSION SCREEN

Depression is measured using the Patient Health Questionnaire-9 (PHQ-9), a nine-item instrument based on the symptoms provided in the Diagnostic and Statistical Manual for Mental Disorders, 4th edition for a major depressive episode in the past two weeks (Spitzer, Kroenke, & Williams, 1999). Following the standard algorithm for interpreting the PHQ-9, symptom levels are categorized as major depression, other depression (less severe depression such as dysthymia or depression not otherwise specified), or neither.


Major depression



Other depression




Any depression




ANXIETY SCREEN

Anxiety is measured using the GAD-7, a seven-item screening tool for screening and severity measuring of generalized anxiety disorder in the past two weeks (Spitzer, Kroenke, Williams, & Lowe, 2006). Following the standard algorithm for interpreting the GAD-7, symptom levels are categorized as severe anxiety, moderate anxiety, or neither.


Severe anxiety



Moderate anxiety




Any anxiety



EATING DISORDER SCREEN


Eating disorders are measured using the written U.S. version of the SCOFF, a five-item screening tool designed to identify subjects likely to have an eating disorder (Morgan, Reid, & Lacey, 1999).

Eating disorders




SUICIDALITY AND SELF-INJUROUS BEHAVIOR


Suicidal ideation (past year)




Suicide plan (past year)



Suicide attempt (past year)



Non-suicidal self-injury (past year)




LIFETIME DIAGNOSES OF MENTAL DISORDERS

Have you ever been diagnosed with any of the following conditions by a health professional (e.g. primary care doctor, psychiatrist, psychologist, etc.)? (Select all that apply)

20%	Depression or other mood disorders (e.g., major depressive disorder, bipolar/manic depression, dysthymia)
21%	Anxiety (e.g., generalized anxiety disorder, phobias, obsessive-compulsive disorder, post-traumatic stress disorder)
10%	Attention disorder or learning disability (e.g., attention deficit disorder, attention deficit hyperactivity disorder, learning disability)
3%	Eating disorder (e.g., anorexia nervosa, bulimia nervosa)
1%	Psychosis (e.g., schizophrenia, schizo-affective disorder)
1%	Personality disorder (e.g., antisocial personality disorder, paranoid personality disorder, schizoid personality disorder)
2%	Substance abuse disorder (e.g., alcohol abuse, abuse of other drugs)
66%	No, none of these


ACADEMIC IMPAIRMENT

In the past 4 weeks, how many days have you felt that emotional or mental difficulties have hurt your academic performance?



POSITIVE MENTAL HEALTH

Positive mental health



Positive mental health (psychological well-being) is measured using The Flourishing Scale, an eight-item summary measure of the respondent's self-perceived success in important areas such as relationships, self-esteem, purpose, and optimism (Diener, Wirtz, Tov, Kim-Prieto, Choi, Oishi, & Biswas-Diener, 2009). The score ranges from 8-56, and we are using 48 as the threshold for positive mental health. In the national 2014 HMS sample, approximately 50% of students are at this threshold or higher.

HEALTH BEHAVIORS AND LIFESTYLE

How often, if ever, have you used any of the substances listed below? Do not include anything you used prescribed to you by a doctor.

Drug use


Never used	Used, but NOT in past 12 months	Used in past 12 months, but NOT in past 30 days	Used in past 30 days	
62%	14%	8%	16%	Cigarettes
53%	15%	13%	19%	Marijuana/cannabis (hashish, blunts)
91%	5%	3%	2%	Cocaine (crack, coke)
94%	3%	1%	1%	Barbiturates or sedatives (prescription-type sleeping pills like Seconal, Ambien, Nembutal, downs or Yellow Jackets)
92%	4%	2%	2%	Tranquilizers (prescription-type drugs like Valium, Librium, Xanax, Ativan, Klonopin)
88%	5%	4%	4%	Amphetamines (Adderall, Ritalin, methamphetamines, crystal meth, speed, uppers, ups)
99%	1%	0%	0%	Heroin
85%	7%	5%	3%	Pain relievers/other opiate-type prescription drugs (codeine, morphine, Demerol, Percodan, Percocet, Vicodin, Oxycontin/oxycodon)
92%	4%	3%	1%	LSD
89%	7%	4%	1%	Other psychedelics or hallucinogens like mushrooms, mescaline or PCP
90%	6%	4%	1%	Ecstasy (MDMA)
98%	1%	0%	0%	Club drugs (Special K, Super K, Ketamine, Liquid G, GHB)
65%	13%	14%	8%	Waterpipe smoking (hookah, arghile, shisha)

Binge drinking

The following questions ask about how much you drink. A "drink" means any of the following:


- A 12-ounce can or bottle of beer
- A 4-ounce glass of wine
- A shot of liquor straight or in a mixed drink

During the last two weeks, how many times have you had 4 (female), 5 (male), 4 or 5 (other gender) or more drinks in a row?



Exercise

In the past 30 days, about how many hours per week on average did you spend exercising? (include any exercise of moderate or higher intensity, where "moderate intensity" would be roughly equivalent to brisk walking or bicycling)




ATTITUDES AND BELIEFS ABOUT MENTAL HEALTH SERVICES


ATTITUDES

Perceived public stigma


Most people would willingly accept someone who has received mental health treatment as a close friend.



Most people feel that receiving mental health treatment is a sign of personal failure.




Most people think less of a person who has received mental health treatment.




Personal stigma


I would willingly accept someone who has received mental health treatment as a close friend.



I feel that receiving mental health treatment is a sign of personal failure.




I would think less of a person who has received mental health treatment.



KNOWLEDGE

Knowledge of campus mental health resources

If you needed to seek professional help for your mental or emotional health while attending, you would know where to go.



USE OF SERVICES

Psychotropic medication use, all students (past year)

In the past 12 months have you taken any of the following types of medications? Please count only those you took, or are taking, several times per week. (Select all that apply)

5%	Psychostimulants (e.g., methylphenidate (Ritalin, or Concerta), amphetamine salts (Adderall), dextroamphetamine (Dexedrine), etc.)
11%	Antidepressants (e.g., fluoxetine (Prozac), sertraline (Zoloft), paroxetine (Paxil), escitalopram (Lexapro), venlafaxine (Effexor), bupropion (Wellbutrin), etc.)
1%	Anti-psychotics (e.g., haloperidol (Haldol), clozapine (Clozaril), risperidone (Risperdal), olanzapine (Zyprexa), etc.)
7%	Anti-anxiety medications (e.g., lorazepam (Ativan), clonazepam (Klonopin), alprazolam (Xanax), buspirone (BuSpar), etc.)
2%	Mood stabilizers (e.g., lithium, valproate (Depakote), lamotrigine (Lamictal), carbamazepine (Tegretol), etc.)
4%	Sleep medications (e.g., zolpidem (Ambien), zaleplon (Sonata), etc.)
2%	Other medication for mental or emotional health
82%	None


Psychotropic medication use among students with positive depression or anxiety screens (past year)

In the past 12 months have you taken any of the following types of medications? Please count only those you took, or are taking, several times per week. (Select all that apply)

8%	Psychostimulants (e.g., methylphenidate (Ritalin, or Concerta), amphetamine salts (Adderall), dextroamphetamine (Dexedrine), etc.)
19%	Antidepressants (e.g., fluoxetine (Prozac), sertraline (Zoloft), paroxetine (Paxil), escitalopram (Lexapro), venlafaxine (Effexor), bupropion (Wellbutrin), etc.)
2%	Anti-psychotics (e.g., haloperidol (Haldol), clozapine (Clozaril), risperidone (Risperdal), olanzapine (Zyprexa), etc.)
12%	Anti-anxiety medications (e.g., lorazepam (Ativan), clonazepam (Klonopin), alprazolam (Xanax), buspirone (BuSpar), etc.)
3%	Mood stabilizers (e.g., lithium, valproate (Depakote), lamotrigine (Lamictal), carbamazepine (Tegretol), etc.)
7%	Sleep medications (e.g., zolpidem (Ambien), zaleplon (Sonata), etc.)
3%	Other medication for mental or emotional health
70%	None


Mental health counseling/therapy, all students (past year)

In the past 12 months have you received counseling or therapy for your mental or emotional health from a health professional (such as psychiatrist, psychologist, social worker, or primary care doctor)?



Mental health counseling/therapy among students with positive depression or anxiety screens (past year)

In the past 12 months have you received counseling or therapy for your mental or emotional health from a health professional (such as psychiatrist, psychologist, social worker, or primary care doctor)?



Informal help-seeking

In the past 12 months have you received counseling or support for your mental or emotional health from any of the following sources? (Select all that apply)

21%	Roommate
50%	Friend (who is not a roommate)
32%	Significant other
42%	Family member
5%	Religious counselor or other religious contact
2%	Support group
2%	Other non-clinical source
32%	None of the above

Barriers to help-seeking


In the past 12 months, which of the following factors have caused you to receive fewer services (counseling, therapy, or medications) for your mental or emotional health than you would have otherwise received? (Select all that apply)

22%	There are financial reasons (too expensive, no insurance)
7%	The location is inconvenient
11%	The hours are inconvenient
35%	I don't have enough time
5%	The number of sessions is too limited
6%	The waiting time until I can get an appointment is too long
11%	I am concerned about privacy
13%	I worry about what others will think of me
7%	I worry that my actions will be documented in my academic record
9%	I worry that my actions will be documented in my medical record
8%	I worry that someone will notify my parents
7%	I fear being hospitalized
3%	People providing services aren't sensitive enough to cultural issues
2%	People providing services aren't sensitive enough to sexual identity issues
1%	I have a hard time communicating in English
10%	I question the quality of my options
18%	I question whether medication or therapy is helpful
8%	I have had bad experiences with medication and/or therapy
18%	The problem will get better by itself
33%	I question how serious my needs are
9%	I don't think anyone can understand my problems
39%	Stress is normal in college/graduate school
29%	I get a lot of support from other sources, such as friends and family
42%	I prefer to deal with issues on my own
5%	Other
9%	There have been no barriers that I can think of


ACADEMIC AND SOCIAL ENVIRONMENT

Campus climate


At my school, I feel that students' mental and emotional well-being is a priority.



At my school, I feel that the academic environment has a negative impact on students' mental and emotional well-being.




At my school, I feel that the campus climate encourages free and open discussion about mental and emotional health.




Support from family

I get the emotional help and support I need from my family.



Support from friends

My friends really try to help me.



BYSTANDER INTERVENTION

Intervening

In the past year, I have intervened in the following situations: (Select all that apply)

32%	Someone was drinking too much
7%	Someone was at risk of being sexually assaulted
32%	Someone was using hurtful language (e.g., bullying, sexist, racist, or homophobic comments)
38%	Someone was experiencing significant emotional distress or thoughts of suicide
3%	Other
37%	None of the above

Witnessing but not intervening

In the past year, I witnessed the following risky or difficult situations but did NOT intervene: (Select all that apply)

23%	Someone was drinking too much
2%	Someone was at risk of being sexually assaulted
16%	Someone was using hurtful language (e.g., bullying, sexist, racist, or homophobic comments)
6%	Someone was experiencing significant emotional distress or thoughts of suicide
1%	Other
65%	None of the above

Reasons for not intervening

In cases where I decided not to intervene, this was because: (Select all that apply)

12%	I was afraid of embarrassing myself
16%	I assumed someone else would do something
38%	I didn't know what to do
29%	I didn't feel confident
58%	I felt it was none of my business
7%	I was afraid my friends wouldn't support me
21%	I felt it was unsafe
14%	I was afraid I'd get in trouble
6%	Other

REFERENCES

MENTAL HEALTH SCREENS

- Diener, E., Wirtz, D., Tov, W., Kim-Prieto, C., Choi, D., Oishi, S., & Biswas-Diener, R. (2009). New measures of well-being: Flourishing and positive and negative feelings. *Social Indicators Research*, 39, 247-266.
- Morgan, J. F., Reid, F., & Lacey, J. H. (1999). The SCOFF questionnaire: assessment of a new screening tool for eating disorders *BMJ*, 319(7223), 1467-1468.
- Spitzer, R. L., Kroenke, K., Williams, J. B., & Patient Health Questionnaire Primary Care Study Group. (1999). Validation and utility of a self-report version of PRIME-MD: the PHQ primary care study. *JAMA*, 282(18), 1737-1744.
- Spitzer, R. L., Kroenke, K., Williams, J. B., & Löwe, B. (2006). A brief measure for assessing generalized anxiety disorder: the GAD-7. *Archives of Internal Medicine*, 166(10), 1092-1097.

SELECTED ARTICLES PUBLISHED WITH HMS DATA

- Eisenberg, D., Golberstein, E., Hunt, J. (2009). Mental Health and Academic Success in College. *B.E. Journal of Economic Analysis & Policy* 9(1) (Contributions): Article 40.
- Eisenberg, D., Hunt, J.B., Speer, N., Zivin, K. (2011). Mental Health Service Utilization among College Students in the United States. *Journal of Nervous and Mental Disease* 199(5): 301-308.
- Eisenberg, D., Chung, H. (2012). Adequacy of Depression Treatment in College Student Populations. *General Hospital Psychiatry* 34(3):213-220.
- Eisenberg, D., Speer, N., Hunt, J.B. (2012). Attitudes and Beliefs about Treatment among College Students with Untreated Mental Health Problems. *Psychiatric Services* 63(7): 711-713.
- Eisenberg, D., Hunt, J.B., Speer, N. (2013). Mental Health in American Colleges and Universities: Variation across Student Subgroups and across Campuses. *Journal of Nervous and Mental Disease* 201(1): 60-67.

Email: healthyminds@umich.edu
Website: www.healthymindsnetwork.org



APPENDIX: DESCRIPTIVE STATISTICS FOR ALL SURVEY ITEMS

MEASURE

All Students

95% CONFIDENCE INTERVAL

Respondent Characteristics

Sample		
N	16342	
Response Rate	29%	
Gender		
Female	55%	(54%, 56%)
Male	45%	(44%, 46%)
Transgender	1%	(1%, 1%)
Race/Ethnicity		
White / Caucasian	73%	(72%, 74%)
Black / African American	6%	(5%, 6%)
Hispanic / Latino	9%	(8%, 10%)
American Indian	1%	(1%, 1%)
Arab / Middle Eastern	1%	(1%, 1%)
Asian / Asian American	13%	(12%, 13%)
Pacific Islander	1%	(0%, 1%)
Other	2%	(2%, 3%)
Country		
US Resident / Citizen	92%	(92%, 93%)
International	8%	(7%, 8%)
Residence		
Campus residence hall	27%	(26%, 28%)
Fraternity / sorority house	1%	(1%, 1%)
Other campus housing	4%	(4%, 4%)
Off-campus / non-university housing	52%	(51%, 53%)
Parent or guardian's home	14%	(13%, 14%)
Other	2%	(2%, 3%)
Academic level		
Associates	14%	(13%, 15%)
Bachelors	71%	(70%, 72%)
Masters	8%	(8%, 9%)
JD	0%	(0%, 0%)
MD	1%	(0%, 1%)
PhD or equivalent	5%	(5%, 5%)
Other	3%	(2%, 3%)
Ever trained for or served in the military (Armed Forces, Reserves, or National Guard)	2%	(2%, 3%)
Age		
18-22	67%	(67%, 68%)
23-25	13%	(12%, 13%)
26-30	9%	(9%, 10%)
31+	11%	(10%, 11%)
Highest educational attainment of either parent		
Less than high school degree	4%	(3%, 4%)
High school degree	24%	(24%, 25%)
College degree	37%	(36%, 38%)
Graduate degree	35%	(34%, 36%)
Religiosity		
Very religious	9%	(8%, 9%)
Fairly religious	24%	(23%, 24%)
Not too religious	28%	(28%, 29%)
Not religious at all	39%	(38%, 40%)

MEASURE

All Students

95% CONFIDENCE INTERVAL

Current financial situation		
It's a financial struggle	22%	(21%, 23%)
It's tight, but I'm doing fine	54%	(53%, 55%)
Finances aren't really a problem	24%	(23%, 24%)
Financial situation growing up		
Very poor, not enough to get by	5%	(4%, 5%)
Enough but not many extras	35%	(34%, 36%)
Comfortable	49%	(48%, 50%)
Well to do	11%	(11%, 12%)
Relationship status		
Single	51%	(50%, 52%)
In a relationship	37%	(36%, 38%)
Married or domestic partnership	11%	(10%, 11%)
Divorced	1%	(1%, 2%)
Sexual orientation		
Heterosexual	84%	(83%, 85%)
Bisexual	5%	(5%, 6%)
Gay / lesbian	5%	(5%, 6%)
Questioning	2%	(2%, 3%)
Other	3%	(3%, 3%)

Mental Health Measures

Positive Mental Health (Diener et al., 2009)		
Flourishing Scale	46.2	(46.0, 46.3)
Depression (PHQ-9)		
Overall score (0-27)	7.0	(6.9, 7.1)
In moderate range (10-14)	15%	(15%, 16%)
In moderately severe range (15-19)	7%	(7%, 8%)
In severe range (20-27)	4%	(4%, 5%)
Major depression (positive screen)	12%	(12%, 13%)
Other depression (positive screen)	9%	(8%, 9%)
Depression overall	21%	(20%, 22%)
Impairment from depression(1)		
Not difficult at all	34%	(33%, 35%)
Somewhat difficult	52%	(51%, 53%)
Very difficult	10%	(10%, 11%)
Extremely difficult	3%	(3%, 3%)
Generalized anxiety (GAD-7)		
Overall score (0-21)	5.9	(5.8, 1.0)
In moderate range (10-14)	13%	(13%, 14%)
In severe range (15-21)	8%	(8%, 9%)
Probable anxiety disorder (positive screen)	22%	(21%, 23%)
Depression/Anxiety		
Depression or anxiety disorder	31%	(30%, 32%)
Disordered eating and body image		
Probable eating disorder (3+ on SCOFF)	7%	(7%, 8%)
Need to be very thin to feel good about self	18%	(17%, 19%)
Think you are very underweight	1%	(1%, 1%)
Academic impairment from mental health, past 4 weeks(2)		
None	40%	(39%, 41%)
1-2 days	28%	(28%, 29%)
3-5 days	17%	(17%, 18%)
6 or more days	14%	(13%, 15%)

(1) How difficult have these problems made it for you to do your work, take care of things at home, or get along with other people?

(2) How many days have you felt that emotional or mental difficulties have hurt your academic performance?

Self-Injury and Suicide

Non-suicidal self-injury, past year		
Any (total)	20%	(19%, 21%)
Cutting self	5%	(4%, 5%)
Burning self	2%	(1%, 2%)
Punching or banging self	6%	(6%, 7%)
Scratching self	7%	(6%, 7%)
Pulling one's hair	6%	(5%, 6%)
Biting self	4%	(4%, 5%)
Interfering with wound healing	5%	(5%, 6%)
Carving words or symbols in skin	1%	(1%, 1%)
Rubbing sharp objects on skin	2%	(2%, 2%)
Punching or banging wall or object	6%	(6%, 7%)
Other	2%	(2%, 2%)
Frequency of self-injury, past year (among those with any)		
Once or twice	55%	(53%, 57%)
Once a month or less	23%	(21%, 25%)
2 or 3 times a month	13%	(11%, 14%)
Once or twice a week	4%	(3%, 5%)
3 to 5 days a week	2%	(2%, 3%)
Nearly everyday, or everyday	2%	(2%, 3%)
Suicidality		
Seriously thought about attempting suicide, past year	10%	(10%, 11%)
Made a plan for attempting suicide, past year	3%	(3%, 4%)
Attempted suicide, past year	1%	(1%, 1%)

Previous Diagnoses of Mental Disorders

Mental disorders		
Any	34%	(33%, 35%)
Depression or mood disorder		
Any	20%	(19%, 21%)
Major depression	6%	(6%, 7%)
Dysthymia	4%	(3%, 4%)
Bipolar	3%	(2%, 3%)
Cyclothymia	1%	(1%, 1%)
Anxiety disorder		
Any	21%	(20%, 21%)
Generalized anxiety disorder	14%	(13%, 14%)
Panic disorder	4%	(3%, 4%)
Agoraphobia	0%	(0%, 0%)
Specific phobia	1%	(1%, 1%)
Social phobia	2%	(2%, 2%)
Obsessive-compulsive disorder (OCD)	3%	(2%, 3%)
Acute stress disorder	1%	(1%, 1%)
Post traumatic stress disorder (PTSD)	3%	(2%, 3%)
Attention or learning disorder		
Any	10%	(9%, 10%)
ADHD	9%	(8%, 9%)
Learning disorder	1%	(1%, 1%)
Eating disorder		
Any	3%	(3%, 3%)
Anorexia nervosa	2%	(1%, 2%)
Bulimia	1%	(1%, 1%)
Binge eating disorder	1%	(0%, 1%)

Previous Diagnoses of Mental Disorders

Psychotic disorder		
Any	1%	(0%, 1%)
Schizophrenia	0%	(0%, 0%)
Personality disorder		
Any	1%	(1%, 2%)
Substance abuse disorder		
Any	2%	(1%, 2%)
Alcohol abuse disorder	1%	(1%, 1%)

Health Behaviors and Lifestyle

Binge drinking, past 2 weeks		
1 or more time	41%	(40%, 41%)
3 or more times	14%	(13%, 14%)
Substance use, past 30 days		
Cigarettes	16%	(15%, 17%)
Marijuana	19%	(18%, 20%)
Cocaine	1%	(0%, 1%)
Cocaine	2%	(1%, 2%)
Barbiturates or sedatives	1%	(1%, 1%)
Tranquilizers	2%	(2%, 2%)
Amphetamines	4%	(4%, 4%)
Heroin	0%	(0%, 0%)
Other opiate-type prescription drugs	3%	(2%, 3%)
LSD	1%	(1%, 1%)
Other psychedelics or hallucinogens	1%	(1%, 1%)
Ecstasy	1%	(1%, 1%)
Club drugs	0%	(0%, 0%)
Waterpipe smoking	8%	(8%, 9%)
Gambled, past year		
Any	15%	(14%, 15%)
On 10+ days	4%	(4%, 5%)
Exercised, past month		
3 or more hours per week	50%	(49%, 51%)
Obese (BMI\geq30)	14%	(13%, 15%)
School work, hours per day in current semester	4.7	(4.6, 4.7)
Sleep (typical amounts during school year)		
Weeknights, hours per night	7.4	(7.4, 7.5)
Weekend nights, hours per night	8.4	(8.4, 8.4)
Naps, hours per day	0.3	(0.3, 0.3)

Attitudes and Beliefs about Services

Stigma about mental health treatment Perceived public stigma score (0-5 scale) Personal stigma score (0-5 scale)	2.1 0.8	(2.0, 2.1) (0.8, 0.8)
...would accept someone who has received MH treatment as a close friend. I... Most people...	98% 85%	(97%, 98%) (85%, 86%)
...think less of someone who has received MH treatment. I... Most people...	7% 47%	(7%, 8%) (47%, 48%)
...feel that receiving mental health treatment is a sign of personal failure. I... Most people...	15% 52%	(14%, 16%) (51%, 53%)
Knows where to go for professional help for mental health Agree or strongly agree	65%	(64%, 66%)
Beliefs about effectiveness of treatment for depression Believes medication is quite helpful or very helpful for depression Believes therapy is quite helpful or very helpful for depression	48% 71%	(47%, 49%) (70%, 72%)
Heard from other students about campus mental health services Mostly negative opinions Even mix of negative and positive opinions Mostly positive opinions Haven't heard anything	5% 12% 20% 63%	(4%, 5%) (12%, 13%) (19%, 21%) (62%, 64%)
Number of close friends or family who have received mental health treatment None 1-2 3+ Don't know	15% 42% 35% 8%	(14%, 15%) (41%, 43%) (34%, 36%) (8%, 9%)

Help Seeking

<i>Think you needed help for emotional or mental health problems, past year</i>		
Strongly agree	19%	(18%, 20%)
Agree	22%	(21%, 22%)
Neither agree nor disagree	14%	(13%, 14%)
Disagree	19%	(18%, 20%)
Strongly disagree	27%	(26%, 28%)
<i>Psychotropic medication</i>		
Any, past year	18%	(17%, 19%)
Any, current	13%	(13%, 14%)
Psychostimulants	5%	(5%, 6%)
Antidepressants	11%	(10%, 11%)
Antipsychotics	1%	(1%, 1%)
Antianxiety	7%	(6%, 7%)
Mood stabilizers	2%	(1%, 2%)
Other	2%	(2%, 2%)
<i>Prescriber (among those with any past-year medication use):</i>		
General practitioner/nurse practitioner/primary care physician	49%	(46%, 51%)
Psychiatrist	38%	(36%, 40%)
Other type of health provider	4%	(3%, 5%)
No prescription	14%	(12%, 16%)
Don't know	2%	(1%, 2%)
<i>Discussed medication with provider, past year (among those with medication use):</i>		
Not at all	18%	(16%, 20%)
1-2 times	42%	(40%, 45%)
3-5 times	21%	(19%, 23%)
More than 5 times	17%	(16%, 19%)
<i>Whom you would talk to, if you were experiencing serious emotional distress:</i>		
Professional clinician	43%	(42%, 44%)
Roommate	24%	(23%, 24%)
Friend (who is not a roommate)	62%	(61%, 63%)
Significant other	45%	(44%, 46%)
Family member	63%	(62%, 64%)
Religious counselor / other religious contact	10%	(9%, 10%)
Support group	4%	(3%, 4%)
Other non-clinical source	2%	(1%, 2%)
No one	6%	(5%, 6%)
<i>Therapy or counseling for mental health</i>		
Past year	22%	(21%, 22%)
Current	10%	(9%, 10%)
<i>Visits in past year, among those with any</i>		
1-3	35%	(33%, 37%)
4-6	25%	(23%, 27%)
7-9	13%	(12%, 15%)
More than 10	27%	(25%, 29%)

Help Seeking

<i>Use of specific providers for therapy or counseling for mental health</i>		
Psychiatric emergency services	0%	(0%, 1%)
Inpatient psychiatric hospital	0%	(0%, 1%)
Partial hospitalization program	0%	(0%, 0%)
Provider in the local community (not on campus)	5%	(5%, 6%)
Provider in another location (such as hometown)	6%	(5%, 6%)
Other	2%	(1%, 2%)
<i>Any medication or therapy for mental health</i>		
Past year	30%	(29%, 31%)
Current	18%	(18%, 19%)
<i>Any medication or therapy, among those with positive depression or anxiety screen</i>		
Past year	46%	(45%, 48%)
Current	30%	(28%, 32%)
<i>Any visit to a health provider</i>		
Past year	79%	(78%, 80%)
<i>Received counseling or support for mental health from these sources, past year:</i>		
Roommate	21%	(21%, 22%)
Friend (other than roommate)	50%	(49%, 51%)
Significant other	32%	(31%, 33%)
Family member	42%	(41%, 43%)
Religious contact	5%	(5%, 6%)
Support group	2%	(2%, 2%)
Other non-clinical source	2%	(2%, 2%)
None of the above	32%	(31%, 33%)

Bystander ("Upstander") Intervention

<i>Intervened in the following situations (past year)</i>		
Someone was drinking too much	32%	(31%, 33%)
Someone was at risk of being sexually assaulted	7%	(7%, 8%)
Someone was using hurtful language (e.g., bullying, sexist, racist, homophobic)	32%	(32%, 33%)
Someone was experiencing significant emotional distress or thoughts of suicide	38%	(37%, 39%)
Other	3%	(3%, 4%)
None of the above	37%	(37%, 38%)
<i>Witnessed the following situations but did NOT intervene (past year)</i>		
Someone was drinking too much	23%	(22%, 24%)
Someone was at risk of being sexually assaulted	2%	(2%, 2%)
Someone was using hurtful language (e.g., bullying, sexist, racist, homophobic)	16%	(15%, 17%)
Someone was experiencing significant emotional distress or thoughts of suicide	6%	(5%, 6%)
Other	1%	(1%, 1%)
None of the above	65%	(64%, 66%)

MEASURE

All Students

95% CONFIDENCE INTERVAL

<i>In cases where I decided not to intervene, this was because:</i>		
I was afraid of embarrassing myself	12%	(11%, 13%)
I assumed someone else would do something	16%	(15%, 17%)
I didn't know what to do	38%	(36%, 39%)
I didn't feel confident	29%	(27%, 30%)
I felt it was none of my business	58%	(56%, 59%)
I was afraid my friends wouldn't support me	7%	(6%, 8%)
I felt it was unsafe	21%	(20%, 22%)
I was afraid I'd get in trouble	6%	(5%, 7%)
Other	14%	(13%, 15%)

Satisfaction with Therapy, Campus Providers

<i>Convenient hours</i>		
Very dissatisfied	1%	(1%, 1%)
Dissatisfied	2%	(1%, 2%)
Somewhat dissatisfied	3%	(2%, 3%)
Somewhat satisfied	6%	(5%, 6%)
Satisfied	14%	(13%, 15%)
Very satisfied	7%	(6%, 8%)
<i>Location</i>		
Very dissatisfied	0%	(0%, 1%)
Dissatisfied	1%	(0%, 1%)
Somewhat dissatisfied	1%	(1%, 1%)
Somewhat satisfied	3%	(2%, 3%)
Satisfied	15%	(14%, 16%)
Very satisfied	13%	(12%, 14%)
<i>Quality of therapists</i>		
Very dissatisfied	1%	(1%, 1%)
Dissatisfied	1%	(1%, 2%)
Somewhat dissatisfied	2%	(2%, 3%)
Somewhat satisfied	5%	(4%, 5%)
Satisfied	12%	(11%, 13%)
Very satisfied	11%	(10%, 12%)
<i>Respect for privacy concerns</i>		
Very dissatisfied	1%	(0%, 1%)
Dissatisfied	0%	(0%, 1%)
Somewhat dissatisfied	1%	(0%, 1%)
Somewhat satisfied	2%	(2%, 2%)
Satisfied	13%	(11%, 14%)
Very satisfied	16%	(15%, 17%)
<i>Scheduling appointments w/o long delays</i>		
Very dissatisfied	1%	(1%, 2%)
Dissatisfied	2%	(1%, 2%)
Somewhat dissatisfied	2%	(2%, 3%)
Somewhat satisfied	5%	(4%, 5%)
Satisfied	12%	(11%, 13%)
Very satisfied	10%	(9%, 11%)

Note: the confidence intervals are wide for these numbers, because the sample sizes are small (these questions were only asked of service users).

Satisfaction with Therapy, Non-Campus Providers

<i>Convenient hours</i>		
Very dissatisfied	3%	(2%, 4%)
Dissatisfied	4%	(2%, 5%)
Somewhat dissatisfied	6%	(5%, 8%)
Somewhat satisfied	15%	(13%, 17%)
Satisfied	44%	(41%, 47%)
Very satisfied	28%	(26%, 31%)
<i>Location</i>		
Very dissatisfied	3%	(2%, 4%)
Dissatisfied	5%	(4%, 6%)
Somewhat dissatisfied	10%	(8%, 11%)
Somewhat satisfied	16%	(14%, 18%)
Satisfied	39%	(36%, 42%)
Very satisfied	27%	(25%, 30%)
<i>Quality of therapists</i>		
Very dissatisfied	4%	(3%, 6%)
Dissatisfied	4%	(3%, 5%)
Somewhat dissatisfied	5%	(3%, 6%)
Somewhat satisfied	10%	(8%, 11%)
Satisfied	31%	(28%, 33%)
Very satisfied	47%	(44%, 49%)
<i>Respect for privacy concerns</i>		
Very dissatisfied	2%	(1%, 3%)
Dissatisfied	1%	(1%, 2%)
Somewhat dissatisfied	2%	(1%, 3%)
Somewhat satisfied	6%	(4%, 7%)
Satisfied	32%	(29%, 35%)
Very satisfied	57%	(54%, 60%)
<i>Scheduling appointments w/o long delays</i>		
Very dissatisfied	3%	(2%, 4%)
Dissatisfied	3%	(2%, 4%)
Somewhat dissatisfied	6%	(5%, 8%)
Somewhat satisfied	11%	(9%, 12%)
Satisfied	35%	(32%, 37%)
Very satisfied	42%	(39%, 44%)

Note: the confidence intervals are wide for these numbers, because the sample sizes are small (these questions were only asked of service users).

Barriers and Facilitators to Help-Seeking

Reasons for receiving no or fewer services for mental health		
No need	43%	(42%, 44%)
Haven't had a chance but plan to go	4%	(4%, 5%)
Prefer to deal with issues on my own	42%	(41%, 43%)
Financial reasons (too expensive, no insurance)	22%	(21%, 23%)
Location is inconvenient	7%	(7%, 8%)
Hours are inconvenient	11%	(10%, 12%)
Don't have enough time	35%	(34%, 36%)
Number of sessions is too limited	5%	(5%, 5%)
Waiting time to get appointment is too long	6%	(6%, 6%)
Concerned about privacy	11%	(10%, 11%)
Worry what others will think	13%	(13%, 14%)
Worry that actions will be documented on academic record	7%	(7%, 8%)
Worry that actions will be documented on medical record	9%	(8%, 9%)
Worry that someone will notify parents	8%	(8%, 9%)
Fear being hospitalized	7%	(6%, 7%)
Providers aren't sensitive enough to cultural issues	3%	(2%, 3%)
Providers aren't sensitive enough to sexual identity issues	2%	(2%, 2%)
Have a hard time communicating in English	1%	(1%, 1%)
Question the quality of options	10%	(10%, 11%)
Question whether medication or therapy is helpful	18%	(17%, 19%)
Have had bad experiences with medication and/or therapy	8%	(7%, 9%)
Problem will get better by itself	18%	(17%, 19%)
Question how serious my needs are	33%	(32%, 34%)
Don't think anyone can understand my problems	9%	(8%, 9%)
Stress is normal in college/graduate school	39%	(38%, 39%)
Get a lot of support from other sources, such as friends and family	29%	(28%, 29%)
Other	5%	(4%, 5%)
None (there have been no barriers than I can think of)	9%	(9%, 10%)
Reasons for seeking help		
Decided on my own	71%	(69%, 73%)
Friend encouraged or pressured me	23%	(21%, 24%)
Family member encouraged or pressured me	37%	(35%, 38%)
Other person encouraged or pressured me	8%	(7%, 10%)
I was mandated by campus staff	2%	(2%, 3%)
I acquired more information about my options	2%	(1%, 2%)
Other reasons	7%	(6%, 8%)

Barriers and Facilitators to Help-Seeking

Source of health insurance		
None (uninsured)	7%	(6%, 7%)
Parent's employer	58%	(57%, 59%)
Own employer	7%	(7%, 8%)
Spouse's employer	3%	(3%, 4%)
Student plan	17%	(16%, 17%)
Embassy or other international source	1%	(0%, 1%)
Individual market	2%	(2%, 2%)
Public insurance	6%	(5%, 6%)
Uncertain whether insured	2%	(1%, 2%)
Insured but uncertain of source	3%	(3%, 3%)
Plan provides any coverage for local MH visits (among those with a plan)		
Yes, it definitely would	24%	(23%, 25%)
I think it would but am not sure	31%	(30%, 32%)
I have no idea	36%	(35%, 37%)
I think it would not but am not sure	7%	(7%, 8%)
No, it definitely would not	2%	(2%, 2%)
Plan meets needs for mental health services (among those with a plan)		
Have not needed plan to cover services	67%	(66%, 67%)
Yes, everything I have needed is covered	26%	(25%, 27%)
No, the coverage is inadequate to meet my needs	7%	(7%, 8%)
Coverage is inadequate because my plan...(among those w/ inadequate plan)		
Doesn't cover any mental health services	19%	(16%, 22%)
Doesn't cover pre-existing conditions	5%	(3%, 7%)
Doesn't cover certain conditions	15%	(12%, 18%)
Has a co-pay that is too expensive	36%	(33%, 40%)
Has a deductible that is too expensive	22%	(19%, 25%)
Doesn't cover certain services or providers	44%	(40%, 48%)
Has a limit on number of services covered	34%	(30%, 37%)

Supportiveness of Academic and Social Environment

Talked with any academic personnel about MH problems affecting performance	16%	(15%, 17%)
Supportiveness of response by academic personnel		
Very supportive	46%	(43%, 48%)
Supportive	44%	(41%, 47%)
Not supportive	8%	(7%, 10%)
Very unsupportive	2%	(2%, 3%)
Whom would you talk to about mental health problems affecting academic performance:		
Professor from one of classes	40%	(40%, 41%)
Academic advisor	32%	(31%, 33%)
Another faculty member	7%	(7%, 8%)
Teaching assistant	3%	(2%, 3%)
Student services staff	14%	(13%, 14%)
Dean of Students or Class Dean	7%	(6%, 7%)
Other	6%	(6%, 7%)
No one	32%	(31%, 33%)

MEASURE

All Students

95% CONFIDENCE INTERVAL

<i>At my school students' mental and emotional well-being is a priority</i>		
Strongly agree	17%	(16%, 18%)
Agree	34%	(33%, 35%)
Neither agree nor disagree	31%	(30%, 32%)
Disagree	13%	(12%, 13%)
Strongly disagree	5%	(5%, 6%)
<i>At my school the academic environment has a negative impact on students' mental and emotional well-being</i>		
Strongly agree	12%	(11%, 13%)
Agree	29%	(28%, 29%)
Neither agree nor disagree	31%	(31%, 32%)
Disagree	22%	(21%, 23%)
Strongly disagree	6%	(5%, 6%)
<i>At my school the campus climate encourages free and open discussion about mental and emotional health</i>		
Strongly agree	8%	(8%, 9%)
Agree	30%	(29%, 30%)
Neither agree nor disagree	41%	(40%, 42%)
Disagree	16%	(15%, 17%)
Strongly disagree	6%	(5%, 6%)
<i>Competitiveness between students in your classes</i>		
Very competitive	18%	(17%, 19%)
Competitive	28%	(27%, 29%)
Somewhat competitive	34%	(33%, 35%)
Not competitive	18%	(17%, 19%)
Very uncompetitive	2%	(2%, 3%)
<i>Receives emotional support from family</i>		
Strongly agree	33%	(32%, 34%)
Agree	27%	(26%, 28%)
Neither agree nor disagree	15%	(15%, 16%)
Disagree	9%	(9%, 10%)
Strongly disagree	15%	(14%, 16%)
<i>Friends really try to help you</i>		
Strongly agree	33%	(32%, 34%)
Agree	32%	(31%, 33%)
Neither agree nor disagree	18%	(17%, 19%)
Disagree	7%	(6%, 7%)
Strongly disagree	10%	(9%, 11%)
<i>Treated unfairly because of race, ethnicity, or culture (past year)</i>		
Never	74%	(73%, 74%)
Once in a while	19%	(18%, 20%)
Sometimes	6%	(5%, 6%)
A lot	1%	(1%, 2%)
Most of the time	0%	(0%, 0%)
Almost all of the time	0%	(0%, 0%)